205.4 - Radioactive Solutions

These SRMs are intended for the calibration of radioactivity measuring instruments and for the monitoring of chemical and geochemical processes. They are calibrated in terms of activity per gram of solution. Each SRM is contained in a flame-sealed glass ampoule or bottle and, except as noted, consists of the radionuclide dissolved in an aqueous solution (usually acidic).

When an import permit for radioactive material is required of a customer outside the U.S., NIST must have a copy to complete an order and facilitate shipment.

"Radioactive SRM Purchasing Instructions & License Certification Form"

PLEASE NOTE: The tables are presented to facilitate comparisons among a family of materials to help customers select the best SRM for their needs. For specific values and uncertainties, the certificate is the only official source.

SRM Description	4222d	4226D	4233f	4239 Strontium-90	4251d	4274 Holmium-166m	4288B	4320b	4321d Natural	4322d	4323c	4324B Uranium-232	4326a	4328C	4329a	4330C
	Carbon-14-n-hexadecane Radioactivity Standard	Nickel-63 Radioactivity Standard	Cesium-137 Radioactivity Standard	Radioactivity	Barium-133 Radioactivity Standard	Gamma-ray	Technetium-99 Radioactivity Standard	Curium-244 Radioactivity Standard	Uranium Radioactivity Standard	Americium-241 Radioactivity Standard	Plutonium-238 Radioactivity Standard	Radioactivity Standard Solution	Polonium-209 Radioactivity Standard	Thorium-229 Radioactivity Standard	Curium-243 Radioactivity Standard	Plutonium-239 Radioactivity Standard
Unit of Issue	(5 mL)	(5 mL)	(5 mL)	(5 mL)	(5 mL)	(5 mL)	(5 mL)	(5 mL)	(5 mL)	(5 mL)	(5 mL)	(5 mL)	(5 mL)	(5 mL)	(5 mL)	(3 mL)
Approx. Massic Activity (Bq/g)	54,000	85 000	221 100	32 000	382 600	20 000	30 000	35	250	134	23	30	39**	35	31	40
Chemical Form	n-hexadecane	NiCl ₂	CsCl	SrCI ₂	BaCl ₂	HoCl3	KTcO ₄	Cm(NO 3)3	UO ₂ (NO ₃) ₂	Am(NO ₃) ₃	Pu(NO ₃) ₆	UO ₂ (NO ₃) ₂	PoCl ₄	Th(NO ₃) ₄	Cm(NO 3)3	Pu(NO ₃) ₆
Decay Modes	β-	β-	β-,γ	β-	EC	Υ	β-	α	α	α	α	α	α, EC	α	α	α
NRC License or Equivalent Required*	-	Х	х	х	х	Х	-	х	х	Х	х	х	х	х	х	Х
Reference Time (month/year	09/14	11/09	08/18	12/06	07/18	02/06	05/08	09/11	03/17	3/19	10/16	7/02	12/13	12/07	05/19	05/09

^{*}If no "X", then license is not required unless the institution possesses a specific license that covers the listed radionuclide.
**Value for Approximate Massic Activity for SRM 4326a is expressed as s⁻¹•g⁻¹.

⁻ Certified values are normal font

Reference values are italicized
 Values in parentheses are for information only

205.4 - Radioactive Solutions

These SRMs are intended for the calibration of radioactivity measuring instruments and for the monitoring of chemical and geochemical processes. They are calibrated in terms of activity per gram of solution. Each SRM is contained in a flame-sealed glass ampoule or bottle and, except as noted, consists of the radionuclide dissolved in an aqueous solution (usually acidic).

When an import permit for radioactive material is required of a customer outside the U.S., NIST must have a copy to complete an order and facilitate shipment.

"Radioactive SRM Purchasing Instructions & License Certification Form"

PLEASE NOTE: The tables are presented to facilitate comparisons among a family of materials to help customers select the best SRM for their needs. For specific values and uncertainties, the certificate is the only official source.

4332E Americium-243 Radioactivity Standard	4334j Plutonium-242 Radioactivity Standard	4337 Lead-210 Radioactivity Standard	4338A Plutonium-240 Radioactivity Standard Solution			
(5 mL)	(5 mL)	(5 mL)	(5 mL)			
40	26	9 000	40			
Am(NO ₃) ₃	Pu(NO ₃) ₆	Pb(NO ₃) ₂	Pu(NO ₃) ₆			
α	α	β-	α			
х	Х	х	Х			
10/08	08/17	06/06	05/96			

^{*}If no "X", then license is not required unless the institution possesses a specific license that covers the listed radionuclide.
**Value for Approximate Massic Activity for SRM 4326a is expressed as s⁻¹•g⁻¹.

⁻ Certified values are normal font

Reference values are italicized
 Values in parentheses are for information only

205.4 - Radioactive Solutions

These SRMs are intended for the calibration of radioactivity measuring instruments and for the monitoring of chemical and geochemical processes. They are calibrated in terms of activity per gram of solution. Each SRM is contained in a flame-sealed glass ampoule or bottle and, except as noted, consists of the radionuclide dissolved in an aqueous solution (usually acidic).

When an import permit for radioactive material is required of a customer outside the U.S., NIST must have a copy to complete an order and facilitate shipment.

"Radioactive SRM Purchasing Instructions & License Certification Form"

PLEASE NOTE: The tables are presented to facilitate comparisons among a family of materials to help customers select the best SRM for their needs. For specific values and uncertainties, the certificate is the only official source.

SRM Description Unit of Issue	4339b Radium-228 Radioactivity Solution (5 mL)	4340B Plutonium-241 Radioactivity Standard (5 mL)	4341a Neptunium-237 Radioactivity Standard (5 mL)	4342A Thorium-230 Radioactivity Standard (5 mL)	4361C Hydrogen-3 Radioactivity Standard (500 mL)	4370d Europium-152 Radioactivity Standard (5 mL)	4915F Cobalt-60 Radioactivity Standard Solution (5 mL)	4919I Strontium-90 Radioactivity Standard (5 mL)	4926E Hydrogen-3 Radioactivity Standard (20 mL)	4927g Hydrogen-3 Radioactivity Standard (5 mL)	4929F Iron-55 Radioactivity Standard (5 mL)	4943 Chlorine-36 Radioactivity Standard (3 mL)	4949d lodine-129 Radioactivity Standard (5 mL)	4965a Radium-226 Radioactivity Standard (5 mL)	4966A Radium-226 Radioactivity Standard (5 mL)	4967A Radium-226 Radioactivity Standard Solution (5 mL)	4969 Radium-226 Radioactivity Standard (5 mL)
Approx. Massic Activity (Bq/g)	200	250	150	40	2	18 700	60 000	4 200	5 000	540 000	59 000	10 000	2 700	30	290	2 500	3
Chemical Form	Ra(NO ₃) ₂	Pu(NO ₃) ₆	Np(NO ₃) ₅	Th(NO ₃) ₄	H ₂ O	EuCl ₃	CoCl ₂	SrCl ₂	H ₂ O	H ₂ O	FeCl3	NaCl	Nal	RaCl ₂	RaCl ₂	RaCl ₂	RaCl ₂
Decay Modes	β-	β-	α	α	β-	β-,EC,γ	β-,γ	β-	β-	β-	EC, β -	β-	β-	α, γ	α, γ	α, γ	α, γ
NRC License or Equivalent Required*	Х	Х	х	х	-	Х	х	х	-	-	-	-	х	х	х	х	х
Reference Time (month/year	10/10	06/07	09/12	04/07	09/98	7/18	11/05	12/06	09/98	05/15	11/05	12/84	01/14	01/07	01/07	09/03	09/98

^{*}If no "X", then license is not required unless the institution possesses a specific license that covers the listed radionuclide.
**Value for Approximate Massic Activity for SRM 4326a is expressed as s⁻¹•g⁻¹.

⁻ Certified values are normal font

Reference values are italicized
 Values in parentheses are for information only